

AUXILIARY SPACE PRECONDITIONERS FOR LINEAR ELASTICITY BASED ON GENERALIZED FINITE ELEMENT METHODS

JAMES BRANNICK AND DURKBIN CHO

ABSTRACT. We construct and analyze a preconditioner of the linear elasticity system discretized by conforming linear finite elements in the framework of the auxiliary space method. The auxiliary space preconditioner is based on discretization of a scalar elliptic equation with Generalized Finite Element Method (GFEM).

1. INTRODUCTION

The discretizations and fast solvers for the linear elasticity systems have been extensive subject of research for the past 20 years. A number of methods have been introduced in [3, 6, 12, 13, 14, 17], and in these works one can find stable discretizations, a priori error estimates, as well as construction of fast and robust solvers for different range of material parameters.

In this paper we consider the variational problem corresponding to the lowest order finite element method discretization of linear elasticity system in displacement formulation. We assume that the material parameters in the linear elasticity system are well behaved, namely, the Poisson ratio is away from $1/2$. We will not discuss the robustness of our method with respect to these parameters, because our goal here is to introduce and prove results on the relationship between GFEM discretizations of scalar equations and the linear elasticity system, as well as to employ such relations in the construction of auxiliary space [15, 18] preconditioner.

The preconditioner that we construct employs as auxiliary space the piecewise quadratic conforming finite elements, and the corresponding auxiliary bilinear form on the auxiliary space corresponds to the discretized scalar Laplace's equation. To relate the discretization of the scalar Laplace problem to linear elasticity, we also use an "intermediate" GFEM space containing piece-wise quadratic functions. The key steps in the analysis of the preconditioner rely on the spectral equivalence results proved in [11] and the fact that the kernel of the quadratic GFEM stiffness matrix is isomorphic to the space of rigid modes (see [11]).

The remainder of the paper is organized as follows. In section 2, we present the linear elasticity problem of interest and its variational formulation. In section 3, we show an auxiliary spectral equivalence relation that plays a key role in the analysis. We introduce the preconditioner, and prove uniform spectral bounds in section 4. Numerical results are presented in section 5 are shown to validate the theoretical results.

Date: March 15, 2010.

2000 Mathematics Subject Classification. 65F10, 65N30, 65N55.

Key words and phrases. linear elasticity, preconditioning, auxiliary space.

2. LINEAR ELASTICITY

Let $\Omega \subset \mathbb{R}^2$ be a bounded convex polygonal domain with boundary $\Gamma = \partial\Omega$. We consider the linear elasticity problem with pure traction boundary conditions:

$$(1) \quad \begin{aligned} -\operatorname{div}(2\mu\varepsilon(\underline{u}) + \lambda\operatorname{tr}(\varepsilon(\underline{u}))\underline{\delta}) &= \underline{f} \quad \text{in } \Omega, \\ (2\mu\varepsilon(\underline{u}) + \lambda\operatorname{tr}(\varepsilon(\underline{u}))\underline{\delta})\underline{n} &= \underline{g} \quad \text{on } \Gamma, \end{aligned}$$

where $\underline{f}$ is an external force, $\underline{n}$ is the outward unit normal on the boundary, $\varepsilon_{ij} = \frac{1}{2}(\partial_j u_i + \partial_i u_j)$ is the strain tensor, and $\underline{\delta}$ is a matrix whose elements consist of the Kronecker delta symbol. Further, let tr denote the trace of a matrix and define the Lamé coefficients μ and λ in terms of the Young modulus E and the Poisson ratio ν as follows:

$$\lambda = \frac{E\nu}{(1+\nu)(1-2\nu)} \quad \text{and} \quad \mu = \frac{E}{2(1+\nu)}$$

We mention that due to the corners of the boundary of the polygonal domain Ω , care must be taken when considering the boundary conditions of (1) (See [6, 8] for details).

Let S_i , $1 \leq i \leq n$, be the vertices of Γ , Γ_i , $1 \leq i \leq n$, the open line segments joining S_i to S_{i+1} , and $\underline{n}_i$ the unit outer normal along Γ_i . Let $p \in H^{1/2}(\Gamma_i)$ and $q \in H^{1/2}(\Gamma_{i+1})$. Then, $p \equiv q$ at S_{i+1} if

$$\int_0^\delta |q(s) - p(-s)|^2 \frac{ds}{s} < \infty,$$

where s is the oriented arc length measured from S_{i+1} , and δ is a positive number less than $\min\{|\Gamma_i| : 1 \leq i \leq n\}$. Then, equation (1) can be written more precisely as

$$(2) \quad \begin{aligned} -\operatorname{div}(2\mu\varepsilon(\underline{u}) + \lambda\operatorname{tr}(\varepsilon(\underline{u}))\underline{\delta}) &= \underline{f} \quad \text{in } \Omega \\ (2\mu\varepsilon(\underline{u}) + \lambda\operatorname{tr}(\varepsilon(\underline{u}))\underline{\delta})\underline{n}_i|_{\Gamma_i} &= \underline{g}_i \quad 1 \leq i \leq n, \end{aligned}$$

where $\underline{f} \in L_2(\Omega)$ and $\underline{g}_i \in H^{1/2}(\Gamma_i)$ satisfy

$$\underline{g}_i \cdot \underline{n}_{i+1} \equiv \underline{g}_{i+1} \cdot \underline{n}_i \quad \text{at } S_{i+1} \quad \text{for } 1 \leq i \leq n.$$

Denote by RM and $\operatorname{RM}^\perp$ the space of rigid modes and its orthogonal complement with respect to $L_2(\Omega)$, respectively and by $H_\perp^1(\Omega) := H^1(\Omega) \cap \operatorname{RM}^\perp$ the intersection of $H^1(\Omega)$ with the L_2 orthogonal complement $\operatorname{RM}^\perp$ of RM . From integration by parts, we obtain the weak formulation of (2) as follows:

Find $\underline{u} \in H_\perp^1(\Omega)$ such that

$$(3) \quad a_{LE}(\underline{u}, \underline{v}) = \int_\Omega \underline{f} \cdot \underline{v} \, dx + \sum_{i=1}^n \int_{\Gamma_i} \underline{g}_i \cdot \underline{v}|_{\Gamma_i} \, ds$$

for all $\underline{v} \in H_\perp^1(\Omega)$, where

$$a_{LE}(\underline{u}, \underline{v}) := \int_\Omega \left(2\mu \langle \varepsilon(\underline{u}), \varepsilon(\underline{v}) \rangle_F + \lambda (\nabla \cdot \underline{u})(\nabla \cdot \underline{v}) \right) dx.$$

Here and below, $\langle \cdot, \cdot \rangle_F$ and $\langle \cdot, \cdot \rangle$ are the Frobenius inner product for matrices and the standard Euclidean inner product for vectors in $\mathbb{R}^d$, respectively. The corresponding norms are denoted with $|\cdot|_F$ and $|\cdot|$, respectively. A sufficient condition for existence of a solution to (3) is that the following compatibility condition is satisfied:

$$\int_{\Omega} \underline{f} \cdot \underline{v} \, dx + \sum_{i=1}^n \int_{\Gamma_i} \underline{g}_i \cdot \underline{v}|_{\Gamma_i} \, ds = 0, \quad \forall \underline{v} \in \text{RM}.$$

Following [19], we write $x_1 \lesssim y_1$, $x_2 \gtrsim y_2$ and $x_3 \bar{\approx} y_3$ whenever there exist constants C_1, c_2, c_3 and C_4 , independent of the mesh size h , such that

$$x_1 \leq C_1 y_1, \quad x_2 \geq c_2 y_2, \quad \text{and} \quad c_3 x_3 \leq y_3 \leq C_4 x_3.$$

Moreover, we write $\lesssim_\lambda, \gtrsim_\lambda$ and $\bar{\approx}_\lambda$ when $\lesssim, \gtrsim$ and $\bar{\approx}$ are dependent on the the Lamé coefficient λ .

3. A MULTILEVEL PRECONDITIONING FOR THE GFEM PROBLEMS

In this section, we present our GFEM-based preconditioner, using results from [11] to motivate our scheme. Let $\Omega \subset \mathbb{R}^d$ be a polygonal domain, with $d = 2, 3$ and V_1 be the associated piecewise linear finite element space on a quasi-uniform triangulation $\mathcal{T}_h$ of Ω . Then the GFEM space V^{GFEM} is defined as

$$V^{GFEM} := V_1 + \sum_{k=1}^d \sum_{i=1}^d \text{span}\{\psi_{k,i}\phi_i\},$$

where V_1 corresponds to continuous piecewise linear elements $\{\phi_i\}_{i=1}^n$ and $\psi_{k,i} = \frac{x_k - x_{k,i}}{h}$ with x_k the k -th component of x and $x_{k,i}$ the k -th value associated with the i -th nodal point. Consider the bilinear form

$$(4) \quad a(u, v) = \int_{\Omega} \nabla u \cdot \nabla v, \quad \forall u, v \in V^{GFEM}.$$

It is shown in [11] that the kernel of the stiffness matrix A corresponding to (4) is characterized by the space RM of rigid modes. To obtain an efficient preconditioner for $a(\cdot, \cdot)$ on V^{GFEM} , we thus choose as auxiliary space $W = V_1 \times V_1^d$ defined by

$$\mathbf{u} \in W, \quad \mathbf{u} = (u, \underline{u}), \quad \underline{u} = (u_k) \quad \text{where} \quad u = \sum_{i=1}^n \alpha_i \phi_i, \quad u_k = \sum_{i=1}^n \alpha_{k,i} \phi_i,$$

equipped with $a_W(\mathbf{u}, \mathbf{v}) = a(u, v) + \int_{\Omega} \langle \varepsilon(\underline{u}), \varepsilon(\underline{v}) \rangle_F \, dx$. Define $\Pi : W \rightarrow V^{GFEM}$ to be:

$$\mathbf{u} \rightarrow \Pi \mathbf{u} = \sum_{i=1}^n \alpha_i \phi_i + \sum_{k=1}^d \sum_{i=1}^n \alpha_{k,i} \psi_{k,i} \phi_i,$$

and take $A_G : V^{GFEM} \rightarrow (V^{GFEM})'$ and $A_W : W \rightarrow W'$ to be the isomorphisms associated with $a(\cdot, \cdot)$ and $a_W(\cdot, \cdot)$, respectively. Here, $'$ and $*$ denote the dual spaces and adjoint operators.

We consider the case when the linear part of the GFEM elements are zero: $W = \{0\} \times V_1^d$, so that $\Pi : W \rightarrow V^{GFEM}$, or more precisely,

$$\mathbf{u} = (0, \underline{u}) \longrightarrow \Pi \mathbf{u} = 0 + \sum_{k=1}^d \sum_{i=1}^n \alpha_{k,i} \psi_{k,i} \phi_i,$$

or

$$\underline{u} \in W \longrightarrow \Pi \underline{u} = \sum_{k=1}^d \sum_{i=1}^n \alpha_{k,i} \psi_{k,i} \phi_i \in V^{GFEM}.$$

We remark that $\ker(A_G)$ is isomorphic to $\ker(A_W)$ where $\ker$ denotes the kernel of the operator. The following spectral equivalence, proved in the next subsection, then holds:

$$(5) \quad a_W(\underline{u}, \underline{u}) \approx a(\Pi \underline{u}, \Pi \underline{u}), \quad \forall \underline{u} \in W.$$

Now, by the auxiliary space lemma [15, 18], $\Pi B \Pi^*$ is a uniform preconditioner for A_G , namely, $\kappa(\Pi B \Pi^* A_G) \lesssim 1$ where B is a BPX preconditioner for A_W (see [11] for a detailed proof).

$$\begin{array}{ccc} W & \xrightarrow{A_W} & W' \\ \Pi \downarrow & & \uparrow \Pi^* \\ V^{GFEM} & \xrightarrow{A_G} & (V^{GFEM})' \end{array}$$

Hereafter, we use Π_G for the GFEM interpolant associated with the GFEM space, V^G .

3.1. Spectral Equivalence. Consider the following bilinear form defined for piecewise linear, continuous vector fields on a triangulation of Ω with simplexes $\mathcal{T}_h$

$$a_W(\underline{u}, \underline{v}) := \int_{\Omega} \langle \varepsilon(\underline{u}), \varepsilon(\underline{v}) \rangle_F.$$

For a given piecewise linear continuous vector field, $\underline{u}$, denote the corresponding element in the GFEM space by u_G . Consider the isomorphism, Π , between the GFEM space (with 0 linear part) and the piece-wise linear continuous vector fields in $\mathbb{R}^d$:

$$u_G = \Pi \underline{u} = \sum_{k=1}^d \sum_{i=1}^n \alpha_{k,i} \psi_{k,i} \phi_i.$$

By direct computation

$$u_G(\mathbf{x}) = \frac{\langle \mathbf{x}, \underline{u} \rangle - \langle \mathbf{x}, \underline{u} \rangle_I}{h},$$

where w_I denotes the continuous linear interpolant of a function w . We now arrive at the following lemma.

Lemma 3.1. *The following relations hold for any fixed simplex T of the triangulation and any $\mathbf{x} \in T$:*

$$(6) \quad u_G(\mathbf{x}) = \frac{1}{h} (\langle \varepsilon(\underline{u}) \mathbf{x}, \mathbf{x} \rangle - \langle \varepsilon(\underline{u}) \mathbf{x}, \mathbf{x} \rangle_I).$$

Proof. The proof follows by taking a Taylor expansion of $\underline{u}$ and using the fact that $\underline{u}$ is a linear vector field on T and hence $\nabla \underline{u}$ is a constant matrix on T :

$$\langle \underline{u}(\mathbf{x}), \mathbf{x} \rangle = \langle \underline{u}(\mathbf{x}_0), \mathbf{x} \rangle + \langle [\nabla \underline{u}] \mathbf{x}, \mathbf{x} \rangle - \langle [\nabla \underline{u}] \mathbf{x}_0, \mathbf{x} \rangle.$$

Taking the linear interpolant on both sides and subtracting leads to

$$u_G(\mathbf{x}) = \frac{1}{h} (\langle [\nabla \underline{u}] \mathbf{x}, \mathbf{x} \rangle - \langle [\nabla \underline{u}] \mathbf{x}, \mathbf{x} \rangle_I).$$

The proof is concluded by observing that for any matrix $Z \in \mathbb{R}^{d \times d}$ we have

$$\langle Z \mathbf{x}, \mathbf{x} \rangle = \frac{1}{2} \langle (Z + Z^T) \mathbf{x}, \mathbf{x} \rangle,$$

where Z^T is the transpose of Z , □

Writing out $1 = \sum_{i=1}^{d+1} \lambda_i$ and $\mathbf{x} = \sum_{i=1}^{d+1} \lambda_i \mathbf{x}_i$, where $\mathbf{x}_i$ are the vertices of T , and $\lambda_i(\mathbf{x})$ are the barycentric coordinate functions, we obtain that

$$(7) \quad u_G = \sum_{E \subset T} \langle \varepsilon(\underline{u}) \frac{(x_i - x_j)}{h}, x_i - x_j \rangle \varphi_E(\mathbf{x})$$

where $\varphi_E = \lambda_i \lambda_j$ on the edge $\overline{x_i x_j}$. Differentiating and taking the L_2 norm we have

$$(8) \quad \|\nabla u_G\|_{0,T}^2 = \sum_{E' \subset T} \sum_{E \subset T} \left(\varepsilon(\underline{u}) \frac{x_E}{h}, x_E \right) \left(\int_T \nabla \varphi_E \cdot \nabla \varphi_{E'} \right) \left(\varepsilon(\underline{u}) \frac{x_{E'}}{h}, x_{E'} \right),$$

where $x_E = x_i - x_j$. The proof of the spectral equivalence lemma uses the following technical result.

Proposition 3.2. *Let $Z \in \mathbb{R}_{sym}^{d \times d}$ be a symmetric matrix and $\tilde{T}$ be a nondegenerate simplex in $\mathbb{R}^d$ whose edges of size (independent on h). Then*

$$|Z|_F^2 \approx \sum_{E \subset \tilde{T}} (\langle Z y_E, y_E \rangle)^2,$$

where the constants of equivalence depend on the spatial dimension d , and y_E is a vector parallel to the edge E .

Proof. We first prove that the following expression is an inner product on $\mathbb{R}_{sym}^{d \times d}$:

$$\langle Y, Z \rangle_* := \sum_{E \subset \tilde{T}} (\langle Z y_E, y_E \rangle \langle Y y_E, y_E \rangle).$$

First, we aim to establish that

$$\langle Z, Z \rangle_* \geq 0, \quad \text{and} \quad \langle Z, Z \rangle_* = 0 \quad \text{iff} \quad Z \equiv 0.$$

We now provide a detailed proof for $d = 3$ case. A similar proof applies to the $d = 2$ case.

Assume that $\langle Z, Z \rangle_* = 0$. This implies that $\langle Z y_E, y_E \rangle = 0$ for each $E \subset \tilde{T}$. Since the size of $\tilde{T}$ is independent of h , we can map it onto the canonical simplex $\hat{T}$ in $\mathbb{R}^d$ (the convex hull of the canonical coordinate vectors in $\mathbb{R}^d$). Clearly such mapping is affine and independent of h as well. Thus, we can limit our proof to $\hat{T}$. Note that three of the edges of the canonical simplex are parallel to the coordinate vectors $\{e_k\}_{k=1}^3$, and there are three more, which are

parallel to the vectors $e_{kl} = (e_k + e_l)$, $1 \leq k < l \leq 3$. Let $Z = \begin{pmatrix} a_1 & a_{12} & a_{13} \\ a_{12} & a_2 & a_{23} \\ a_{13} & a_{23} & a_3 \end{pmatrix}$. From $\langle Ze_k, e_k \rangle = 0$ it follows that $a_k = 0$. Next, $\langle Ze_{kl}, e_{kl} \rangle = 0$ implies that $a_k + a_l + 2a_{kl} = 0$, and hence $a_{kl} = 0$ as well. Proofs of the remaining properties of the inner product are straightforward and hence omitted.

The proof of the proposition is then concluded by using the fact that all norms on the finite dimensional space $\mathbb{R}_{\text{sym}}^{d \times d}$ are equivalent, with equivalence constants depending on the dimension. $\square$

We now prove that the Poisson bilinear form on GFEM and $a_W(\cdot, \cdot)$ are equivalent.

Lemma 3.3. *The following equivalence relation holds*

$$\|\nabla u_G\|_{0,\Omega} \approx \|\varepsilon(u)\|_{0,\Omega}.$$

Proof. Take $T \in \mathcal{T}_h$. Note that u_G is a quadratic function on T , namely

$$u_G = \sum_{E \subset T} \Theta_E \varphi_E, \quad \text{where} \quad \Theta_E = \left(\varepsilon(\underline{u}) \frac{x_E}{h}, x_E \right),$$

This function vanishes at all vertices of T , and its linear interpolant $(u_G)_I$ is zero. Thus,

$$h^{-2} \|u_G\|_{0,T}^2 \lesssim \|\nabla u_G\|_{0,T}^2 \lesssim h^{-2} \|u_G\|_{0,T}^2,$$

where the first inequality follows from the standard interpolation error estimate and the second is an inverse inequality. On the other hand

$$\|u_G\|_{0,T}^2 \approx |T| \sum_{E \subset T} \Theta_E^2.$$

Hence,

$$(9) \quad |\nabla u_G|_T^2 \approx h^{-2} |T| \sum_{E \subset T} \Theta_E^2 = |T| \sum_{E \subset T} \left(\varepsilon(\underline{u}) \frac{x_E}{h}, \frac{x_E}{h} \right)^2.$$

Note that the size of $\frac{x_E}{h}$ is independent of h and thus we can apply Proposition 3.2 (with $\tilde{T}$ homotetic to T , and with edges $y_E = \frac{x_E}{h}$). Hence, from Proposition 3.2 and (9) we obtain

$$\begin{aligned} \|\varepsilon(\underline{u})\|_{0,\Omega}^2 &= \sum_{T \in \mathcal{T}_h} \|\varepsilon(\underline{u})\|_{0,T}^2 = \sum_{T \in \mathcal{T}_h} \int_T \langle \varepsilon(\underline{u}), \varepsilon(\underline{u}) \rangle_F \\ (10) \quad &\approx \sum_{T \in \mathcal{T}_h} |T| \sum_{E \subset T} \left(\varepsilon(\underline{u}) \frac{x_E}{h}, \frac{x_E}{h} \right)^2 \approx \sum_{T \in \mathcal{T}_h} \|\nabla u_G\|_{0,T}^2 \\ &= \|\nabla u_G\|_{0,\Omega}^2. \end{aligned}$$

$\square$

4. PRECONDITIONING FOR LINEAR ELASTICITY

In this section, we develop and analyze an efficient preconditioner for the linear elasticity problem. An additional spectral equivalence is needed to verify the auxiliary space lemma and, hence, the optimality for our preconditioner in this setting. This spectral equivalence result follows from a Korn inequality on $\tilde{H}_\perp^1(\Omega)$ and the Cauchy-Schwarz inequality.

Proposition 4.1 (Korn inequality). *There exists a positive constant C such that*

$$\|\varepsilon(\underline{v})\|_{0,\Omega} \geq C\|\underline{v}\|_{1,\Omega} \quad \forall \underline{v} \in \underline{H}_\perp^1(\Omega).$$

Proof. The proof is given in [1, 7]. □

It follows from the Korn inequality that the weak formulation (3) has a unique solution $\underline{u} \in \underline{H}_\perp^1(\Omega)$.

Theorem 4.2. *Further, the following spectral equivalence holds:*

$$\|\varepsilon(\underline{v})\|_{0,\Omega}^2 \lesssim a_{LE}(\underline{v}, \underline{v}) \lesssim_\lambda \|\varepsilon(\underline{v})\|_{0,\Omega}^2, \quad \forall \underline{v} \in \underline{H}_\perp^1(\Omega).$$

Proof. Let $\underline{v} \in \underline{H}_\perp^1(\Omega)$. Then we have

$$\begin{aligned} a_{LE}(\underline{v}, \underline{v}) &= 2\mu\|\varepsilon(\underline{v})\|_{0,\Omega}^2 + \lambda\|\nabla \cdot \underline{v}\|_{0,\Omega}^2 \\ &\lesssim 2\mu\|\varepsilon(\underline{v})\|_{0,\Omega}^2 + \lambda\|\underline{v}\|_{1,\Omega}^2 \\ &\lesssim 2\mu\|\varepsilon(\underline{v})\|_{0,\Omega}^2 + \lambda\|\varepsilon(\underline{v})\|_{0,\Omega}^2 \\ &\lesssim_\lambda \|\varepsilon(\underline{v})\|_{0,\Omega}^2, \end{aligned}$$

where the Cauchy-Schwarz inequality and the Korn inequality are used. The reverse inequality is verified as follows:

$$\|\varepsilon(\underline{v})\|_{0,\Omega}^2 \lesssim 2\mu\|\varepsilon(\underline{v})\|_{0,\Omega}^2 + \lambda\|\nabla \cdot \underline{v}\|_{0,\Omega}^2 = a_{LE}(\underline{v}, \underline{v}),$$

for $\forall \underline{v} \in \underline{H}_\perp^1(\Omega)$. □

Finally, we proceed to derive our GFEM auxiliary space preconditioner. Consider

$$a_W(\underline{u}, \underline{v}) := \int_\Omega \langle \varepsilon(\underline{u}), \varepsilon(\underline{v}) \rangle_F dx \quad \forall \underline{u}, \underline{v} \in W,$$

and the mapping $\Pi_G : W \rightarrow V^G$ defined as follows: let $\underline{u} \in W$ be such that

$$\underline{u} = (u_k), \quad u_k = \sum_{i=1}^n \alpha_{k,i} \phi_i \quad \text{for } k = 1, \dots, d,$$

then

$$u_G = \Pi_G(\underline{u}) := \sum_{k=1}^2 \sum_{i=1}^n \alpha_{k,i} \psi_{k,i} \phi_i \in V^G.$$

Let V_q denote the space of continuous piecewise quadratic finite elements equipped with standard inner product $a_q(\cdot, \cdot)$ and define the mapping $\Pi_q : V^G \rightarrow V_q$ by

$$\Pi_q u_G := u_G, \quad \text{for } u_G \in V^{GFEM}.$$

We note that Π_q is the natural inclusion from V^{GFEM} into V_q . Associated with $a_q(\cdot, \cdot)$, we write $A_q : V_q \rightarrow (V_q)'$ for its isomorphism. Going back to the original problem (3), we define the conforming finite element spaces on the mesh $\mathcal{T}_h$:

$$\mathcal{V} := \{\underline{u} : \underline{u} = (u_k)_{k=1}^2, u_k \in V_1\}$$

and

$$\mathcal{V}_\perp := \left\{ \underline{u} \in \mathcal{V} : \int_\Omega \underline{u} \cdot \underline{v} \, dx = 0 \quad \forall \underline{v} \in \text{RM} \right\} \subset H_\perp^1(\Omega).$$

We remark that W can be viewed as W/RM since $\text{Ker}(\Pi_G) = \text{RM}$ and then W is isomorphic to $\mathcal{V}_\perp$. To see this, define the operator $Q_{\text{RM}} : W \rightarrow W$ as follows: for $\underline{v} \in W$,

$$(Q_{\text{RM}}\underline{v}, \underline{w})_{0,\Omega} = (\underline{v}, \underline{w})_{0,\Omega}, \quad \underline{w} \in \text{RM} \subsetneq W.$$

Using Q_{RM} , we can regard W as $\text{Range}(I - Q_{\text{RM}})$ where I is an identity operator and Range denotes the range of an operator and, thus, W is isomorphic to $\mathcal{V}_\perp$.

The auxiliary space lemma (See [15, 18] for a detailed proof) for the linear elasticity problem reads:

Lemma 4.3 (Auxiliary Space Lemma). *Assume that $\tilde{\Pi} : V_q \rightarrow W$ is a surjective and bounded linear operator, namely, there exists a positive constant c_1 such that $\forall v_q \in V_q$,*

$$a_W(\tilde{\Pi}v_q, \tilde{\Pi}v_q) \leq c_1 a_q(v_q, v_q).$$

Also, we suppose that there exists a positive constant c_0 such that $\forall \underline{v} \in W$, there is $v_q \in V_q$ so that

$$\underline{v} = \tilde{\Pi}v_q \quad \text{and} \quad a_q(v_q, v_q) \leq c_0 a_W(\underline{v}, \underline{v})$$

Then

$$(11) \quad c_0^{-2} a_W(\underline{u}, \underline{u}) \leq a_W(\tilde{\Pi}A_q^{-1}\tilde{\Pi}^*A_W\underline{u}, \underline{u}) \leq c_1^2 a_W(\underline{u}, \underline{u}), \quad \forall \underline{u} \in W.$$

We recall that $a_W(\cdot, \cdot) \lesssim a_{LE}(\underline{v}, \underline{v}) \lesssim_\lambda a_W(\cdot, \cdot)$. Also, we see that W and $\mathcal{V}_\perp$ are identical. For these reasons, it suffices to construct a preconditioner for $a_W(\cdot, \cdot)$ on the space W . Let Π_q be an inclusion from V^G into V_q . We define its Hilbert adjoint operator $\Pi_q^\dagger : V_q \rightarrow V^G$ by

$$a_q(\Pi_q u_G, v_q) = a_G(u_G, \Pi_q^\dagger v_q), \quad \forall u_G \in V^G, v_q \in V_q.$$

We note that

$$a_G(\Pi_G \underline{u}, \Pi_G \underline{u}) \approx a_W(\underline{u}, \underline{u}), \quad \forall \underline{u} \in W,$$

and

$$a_G(v_G, v_G) \approx a_W(\Pi_G^{-1}v_G, \Pi_G^{-1}v_G), \quad v_G \in V^G,$$

since Π_G is bijective. Also, the Hilbert adjoint operator $(\Pi_G^{-1})^\dagger$ of Π_G^{-1} can be defined as above. Associated with $a_{LE}(\cdot, \cdot)$ in (3), we write its isomorphism $A_{LE} : \mathcal{V}_\perp \rightarrow (\mathcal{V}_\perp)'$. We summarize the relations between these spaces and their associated interpolants and dual space in the commutative diagram provided in Figure 4.

Let $\tilde{\Pi} := \Pi_G^{-1}\Pi_q^\dagger$. Then $\tilde{\Pi}$ is surjective operator since $\Pi_q^\dagger$ is surjective. Letting $v_q \in V_q$, we have

$$\begin{array}{ccc}
V_q & \xrightarrow{A_q} & (V_q)' \\
\Pi_q^\dagger \downarrow & & \uparrow (\Pi_q^\dagger)^* \\
V^G & \xrightarrow{A_G} & (V^G)' \\
\Pi_G^{-1} \downarrow & & \uparrow (\Pi_G^{-1})^* \\
W & \xrightarrow{A_W} & W' \\
\parallel & & \parallel \\
\mathcal{V}_\perp & \xrightarrow{A_{LE}} & (\mathcal{V}_\perp)'
\end{array}$$

FIGURE 1. Commutative diagram for A_{LE} and our auxiliary spaces.

$$\begin{aligned}
a_W(\tilde{\Pi}v_q, \tilde{\Pi}v_q) &= a_W(\Pi_G^{-1}\Pi_q^\dagger v_q, \Pi_G^{-1}\Pi_q^\dagger v_q) \\
&\lesssim a_G(\Pi_q^\dagger v_q, \Pi_q^\dagger v_q) = a_q(\Pi_q \Pi_q^\dagger v_q, v_q) \\
&\leq a_q(\Pi_q \Pi_q^\dagger v_q, \Pi_q \Pi_q^\dagger v_q)^{1/2} a_q(v_q, v_q)^{1/2} \\
&= a_G(\Pi_q^\dagger v_q, \Pi_q^\dagger v_q)^{1/2} a_q(v_q, v_q)^{1/2} \\
&\lesssim a_W(\Pi_G^{-1}\Pi_q^\dagger v_q, \Pi_G^{-1}\Pi_q^\dagger v_q)^{1/2} a_q(v_q, v_q)^{1/2} \\
&= a_W(\tilde{\Pi}v_q, \tilde{\Pi}v_q)^{1/2} a_q(v_q, v_q)^{1/2}.
\end{aligned}$$

Further, let $\underline{v} \in W$ and set $v_q := \Pi_q \Pi_G \underline{v} \in V_q$. Then, for $\forall \underline{u} \in W$, we have

$$\begin{aligned}
a_W(\tilde{\Pi}v_q, \underline{u}) &= a_W(\Pi_G^{-1}\Pi_q^\dagger \Pi_q \Pi_G \underline{v}, \underline{u}) \\
&= a_G(\Pi_q^\dagger \Pi_q \Pi_G \underline{v}, (\Pi_G^{-1})^\dagger \underline{u}) \\
&= a_q(\Pi_q \Pi_G \underline{v}, \Pi_q (\Pi_G^{-1})^\dagger \underline{u}) \\
&= a_G(\Pi_G \underline{v}, (\Pi_G^{-1})^\dagger \underline{u}) \\
&= a_G(\underline{v}, \underline{u}).
\end{aligned}$$

Therefore, $\underline{v} = \tilde{\Pi}v_q$. Moreover,

$$\begin{aligned}
a_q(v_q, v_q) &= a_q(\Pi_q \Pi_G \underline{v}, \Pi_q \Pi_G \underline{v}) \\
&= a_G(\Pi_G \underline{v}, \Pi_G \underline{v}) \\
&\lesssim a_W(\underline{v}, \underline{v}).
\end{aligned}$$

Consequently, we obtain

$$(12) \quad \kappa(\Pi_G^{-1}\Pi_q^\dagger A_q^{-1}(\Pi_q^\dagger)^*(\Pi_G^{-1})^* A_W) = \kappa(\tilde{\Pi} A_q^{-1} \tilde{\Pi}^* A_W) \lesssim 1.$$

For notational brevity, let $B := \tilde{\Pi} A_q^{-1} \tilde{\Pi}^*$. By (12) and Theorem 4.2, we get $\kappa(BA_{LE}) \lesssim \lambda$, independent of mesh size h .

We define the norm $\|\cdot\|_1$ on W as follows:

$$\|\underline{u}\|_1^2 := |u_1|_1^2 + |u_2|_1^2,$$

where $\underline{u} = (u_1, u_2) \in W$ and write A_1 for the isomorphism associated with $\|\cdot\|_1^2$. Then, using the Korn inequality and the Cauchy-Schwarz inequality, we get

$$(13) \quad \int_{\Omega} \langle \varepsilon(\underline{u}), \varepsilon(\underline{u}) \rangle_F dx \approx \|\underline{u}\|_1^2, \quad \forall \underline{u} \in W.$$

These spectral equivalence relations we proposed basically motivate our choice of GFEM-based auxiliary space preconditioner. We also remark that the space $\tilde{H}^1(\Omega)$ has the usual norm

$$\|\underline{u}\|_{\tilde{H}^1(\Omega)} := (\|\underline{u}\|_1^2 + \|\underline{u}\|_{L_2}^2)^{1/2}.$$

We describe an algorithm under which several numerical experiments in next subsection will be performed. Applying the shape functions on FEM into the bilinear forms $a_{LE}(\cdot, \cdot)$, $a_W(\cdot, \cdot)$ and $a_1(\cdot, \cdot)$, we obtain the Galerkin matrix $\mathbf{A}$, which represents matrices $\mathbf{A}_{LE}$, $\mathbf{A}_W$ and $\mathbf{A}_1$. We consider only a linear system

$$\mathbf{A}x = f,$$

where f is appropriately chosen. This simplification makes sense by the spectral equivalent argument we established. The algorithm follows the preconditioned conjugate gradient methods in [16].

Algorithm 4.4.

Compute $r_0 = f - \mathbf{A}x_0$, $z_0 = \tilde{\Pi} \mathbf{B}_q \tilde{\Pi}^T r_0$, and $p_0 = z_0$

For $j = 0, 1, \dots$, until convergence Do:

$$\alpha_j = (r_j, z_j) / (\mathbf{A}p_j, p_j)$$

$$x_{j+1} = x_j + \alpha_j p_j$$

$$r_{j+1} = r_j - \alpha_j \mathbf{A}p_j$$

$$z_{j+1} = \tilde{\Pi} \mathbf{B}_q \tilde{\Pi}^T r_{j+1}$$

$$\beta_j = (r_{j+1}, z_{j+1}) / (r_j, z_j)$$

$$p_{j+1} = z_{j+1} + \beta_j p_j$$

EndDo

Here, $\tilde{\Pi}$ is the matrix representation of $\tilde{\Pi}$ and $\mathbf{B}_q$ is an approximate inverse of $\mathbf{A}_q$ where $\mathbf{A}_q$ is the Galerkin matrix arising from the piecewise quadratic FEM. The numerical experiments in next section are performed using the stopping criterion $\frac{\|r_k\|}{\|r_0\|} < 10^{-8}$, where r_k is the residual of k th iteration and initial guess $x_0 = (1, -1, \dots, 1, -1)^T$. Taking an account into eigenvalues of the preconditioned system, we can estimate the condition number from parameters in the conjugate gradient algorithm (see [16] for more detail).

5. NUMERICAL EXPERIMENTS

Here we report computed estimates of the condition number of our auxiliary-space preconditioned system matrix for the linear elasticity model discretized using various choices

of mesh spacing on unit square $\Omega = [0, 1]^2$. Here, h represents the lengths of the horizontal and vertical sides of triangles in the meshes.




FIGURE 2. Meshes on the unit square

Experiment 1. We consider a pair of poisson equations on Ω with homogeneous Neumann boundaries:

$$(14) \quad \begin{cases} \begin{pmatrix} \Delta & 0 \\ 0 & \Delta \end{pmatrix} \underline{u} = \underline{0} & \text{in } \Omega, \\ \begin{pmatrix} \frac{\partial}{\partial n} & 0 \\ 0 & \frac{\partial}{\partial n} \end{pmatrix} \underline{u} = \underline{0} & \text{on } \partial\Omega. \end{cases}$$

We note that A_1 is the isomorphism associated with the weak formulation of (14) on the space W . The results in Table 1 implies that the condition number of the preconditioned system of A_1 is bounded independent of mesh size h .

h	$\frac{1}{4}$	$\frac{1}{8}$	$\frac{1}{16}$	$\frac{1}{32}$	$\frac{1}{64}$
iter	20	24	27	27	27
κ	1.92e+1	2.80e+1	3.51e+1	4.02e+1	4.39e+1

TABLE 1. Condition numbers $\kappa(\Pi_G^{-1}\Pi_q^\dagger A_q^{-1}(\Pi_q^\dagger)^*(\Pi_G^{-1})^* A_1)$.

Experiment 2. We deal with a special case when $\lambda = 0$. Namely, the linear elasticity reads

$$(15) \quad \begin{aligned} -\operatorname{div}(\varepsilon(\underline{u})) &= \underline{f} \quad \text{in } \Omega \\ \varepsilon(\underline{u})\underline{n}\Big|_{\Gamma_\ell} &= \underline{0}, \quad 1 \leq \ell \leq 4, \end{aligned}$$

where Γ_ℓ represent four sides of the unit square for $\ell = 1, \dots, 4$. Also, without loss of generality, we suppose that $\mu = \frac{1}{2}$, here and later. Assume that the body force $\underline{f} = \begin{pmatrix} f_1 \\ f_2 \end{pmatrix}$ is defined by

$$(16) \quad f_1 = 0 \quad \text{and} \quad f_2 = 0.$$

Then the exact solution $\underline{u} \in \underline{H}_\perp^2(\Omega)$ is

$$\underline{u} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}.$$

We see that A_W is the isomorphism associated with the weak formulation of (15) on W . In Table 2, $\kappa = C/c$, where C and c are the smallest and largest constants satisfying:

$$ca_W(\underline{u}, \underline{u}) \leq (B_q \Pi_q \Pi_G \underline{u}, \Pi_q \Pi_G \underline{u})_0 \leq Ca_W(\underline{u}, \underline{u}), \forall \underline{u} \in \mathcal{V}_\perp.$$

In Table 2, we observe that the preconditioner we devised is an efficient preconditioner for

h	$\frac{1}{4}$	$\frac{1}{8}$	$\frac{1}{16}$	$\frac{1}{32}$	$\frac{1}{64}$
iter	19	19	17	16	15
κ	6.47e+0	6.70e+0	6.71e+0	6.67e+0	6.48e+0

TABLE 2. Condition numbers $\kappa(\Pi_G^{-1} \Pi_q^\dagger A_q^{-1} (\Pi_q^\dagger)^* (\Pi_G^{-1})^* A_W)$.

the special case. Namely, $\kappa(BA_W) \lesssim 1$.

Experiment 3. In this experiment, we study more general linear elasticity problems with various λ :

$$(17) \quad \begin{aligned} -\operatorname{div}(\sigma(\underline{u})) &= \underline{0} \quad \text{in } \Omega = \text{unit square} \\ \sigma(\underline{u}) \underline{n} \Big|_{\Gamma_\ell} &= \underline{0} \quad 1 \leq \ell \leq 4, \end{aligned}$$

where $\sigma(\underline{u}) = \varepsilon(\underline{u}) + \lambda \operatorname{tr}(\varepsilon(\underline{u})) \underline{\delta}$ and Γ_ℓ are defined as in Experiment 2. In Tables 3, 4 and 5, $\kappa = C/c$, where C and c are the smallest and largest constants satisfying:

$$ca_{LE}(\underline{u}, \underline{u}) \leq (B_q \Pi_q \Pi_G \underline{u}, \Pi_q \Pi_G \underline{u})_0 \leq Ca_{LE}(\underline{u}, \underline{u}), \forall \underline{u} \in \mathcal{V}_\perp.$$

h	$\lambda = 1$		$\lambda = 5$		$\lambda = 10$	
	iter	κ	iter	κ	iter	κ
$\frac{1}{4}$	24	1.33e+1	30	4.60e+1	33	8.60e+1
$\frac{1}{8}$	28	1.39e+1	39	4.51e+1	50	8.60e+1
$\frac{1}{16}$	25	1.40e+1	43	4.57e+1	55	8.55e+1
$\frac{1}{32}$	23	1.40e+1	40	4.56e+1	53	8.55e+1
$\frac{1}{64}$	21	1.39e+1	36	4.54e+1	46	8.49e+1

TABLE 3. Condition numbers $\kappa(\Pi_G^{-1} \Pi_q^\dagger A_q^{-1} (\Pi_q^\dagger)^* (\Pi_G^{-1})^* A_{LE})$.

Experiment 4. For small λ , we experiment the linear elasticity with homogeneous pure traction boundary condition:

$$(18) \quad \begin{aligned} -\operatorname{div}(\varepsilon(\underline{u}) + \lambda \operatorname{tr}(\varepsilon(\underline{u})) \underline{\delta}) &= \underline{f} \quad \text{in } \Omega \\ (\varepsilon(\underline{u}) + \lambda \operatorname{tr}(\varepsilon(\underline{u})) \underline{\delta}) \underline{n} \Big|_{\Gamma_\ell} &= \underline{0}_\ell \quad 1 \leq \ell \leq 4, \end{aligned}$$

where Γ_ℓ are also defined as in Experiment 2 and the body force $\underline{f} = \begin{pmatrix} f_1 \\ f_2 \end{pmatrix}$ is such that

$$\underline{u} = \begin{pmatrix} u_1 \\ u_2 \end{pmatrix} = \begin{pmatrix} x(1-x)y^2(1-y)^2 \sin \pi x - \frac{2}{15\pi^3} \\ x^2(1-x)^2 y^2(1-y)^2 \cos \pi y \end{pmatrix} \in \underline{H}_\perp^2(\Omega)$$

is the exact solution of (18). The results reported in Tables 4 indicate that the condition number of our preconditioned operator remains bounded independent of mesh size and, hence, provides for an optimal order preconditioner for our linear elasticity model problem. They coincide with our theoretical result that $\kappa(BA_{LE}) \lesssim \lambda$, independent of h . We note that although our experiments have been carried out on a structured grid, our analysis carries over to unstructured simplicial meshes and our scheme is thus expected to perform similar for unstructured grids as well.

h	$\lambda = 1$			$\lambda = 5$			$\lambda = 10$		
	iter	κ	$\tilde{H}^1$ norm	iter	κ	$\tilde{H}^1$ norm	iter	κ	$\tilde{H}^1$ norm
$\frac{1}{4}$	26	1.42e+1	7.71e-3	32	4.60e+1	1.10e-2	35	8.60e+1	1.25e-2
$\frac{1}{8}$	31	1.39e+1	4.07e-3	46	4.60e+1	7.56e-3	56	8.60e+1	9.62e-3
$\frac{1}{16}$	31	1.41e+1	1.41e-3	53	4.58e+1	3.48e-3	68	8.56e+1	5.21e-3
$\frac{1}{32}$	31	1.41e+1	3.91e-4	54	4.59e+1	1.12e-3	73	8.58e+1	1.92e-3
$\frac{1}{64}$	31	1.41e+1	1.01e-4	54	4.59e+1	3.04e-4	72	8.58e+1	5.49e-4

TABLE 4. Condition numbers $\kappa(\Pi_G^{-1}\Pi_q^\dagger A_q^{-1}(\Pi_q^\dagger)^*(\Pi_G^{-1})^* A_{LE})$ and the discrete $\tilde{H}^1$ errors between the exact solution and the numerical solution.

Challenging experiment. It is well known that the performance of the piecewise linear finite elements deteriorates as λ approaches ∞ . In the elasticity literature, it is called the locking phenomenon (see [5, 4] for more information). To overcome the effects of the locking, several methods have been suggested in [2, 6, 10, 9, 12, 14, 17]. Both our theoretical and numerical results show that the GFEM-based auxiliary preconditioner for the linear elasticity problem with small λ works very efficiently. It would also be worthwhile to investigate the numerical results when the preconditioner is applied to linear elasticity for large λ . We observe in Table 5 that even when λ is large enough, the condition number of the preconditioned systems is bounded independent of mesh size h and the discretization error estimates look reasonable if h is sufficiently small.

h	$\lambda = 50$ ($\nu = 0.49505$)			$\lambda = 100$ ($\nu = 0.49751$)			$\lambda = 500$ ($\nu = 0.49950$)			$\lambda = 1000$ ($\nu = 0.49975$)		
	iter	κ	$\tilde{H}^1$ norm	iter	κ	$\tilde{H}^1$ norm	iter	κ	$\tilde{H}^1$ norm	iter	κ	$\tilde{H}^1$ norm
$\frac{1}{8}$	92	4.06e+2	1.39e-2	109	8.06e+2	1.51e-2	134	4.01e+3	1.65e-2	140	8.01e+3	1.66e-2
$\frac{1}{16}$	114	4.06e+2	1.04e-2	144	8.06e+2	1.25e-2	224	4.01e+3	1.57e-2	258	8.01e+3	1.64e-2
$\frac{1}{32}$	135	4.06e+2	5.75e-3	170	8.06e+2	8.06e-3	287	4.01e+3	1.31e-2	365	8.01e+3	1.47e-2
$\frac{1}{64}$	145	4.05e+2	2.19e-3	183	8.06e+2	3.69e-3	322	4.01e+3	8.78e-3	404	8.01e+3	1.12e-2
$\frac{1}{128}$	143	4.05e+2	6.42e-4	192	8.05e+2	1.21e-3	367	4.00e+3	4.32e-3	455	8.01e+3	6.36e-3

TABLE 5. Condition numbers $\kappa(\Pi_G^{-1}\Pi_q^\dagger A_q^{-1}(\Pi_q^\dagger)^*(\Pi_G^{-1})^* A_{LE})$ and the discrete $\tilde{H}^1$ errors between the exact solution and the numerical solution.

6. CONCLUDING REMARKS

In this paper, using the auxiliary space technique we have designed a preconditioner for the solution of the problem of linear elasticity, which is also based on generalized finite element methods. We have proved that for arbitrarily fixed λ , the GFEM-based auxiliary

preconditioner always works optimally for the system of linear elasticity discretized using the lowest finite element shape functions.

REFERENCES

- [1] Douglas Arnold. Lecture note on linear plate theory: modeling, analysis, and computation.
- [2] Douglas N. Arnold, Franco Brezzi, Richard S. Falk, and L. Donatella Marini. Locking-free Reissner-Mindlin elements without reduced integration. *Comput. Methods Appl. Mech. Engrg.*, 196(37-40):3660–3671, 2007.
- [3] Douglas N. Arnold and Ragnar Winther. Mixed finite elements for elasticity. *Numer. Math.*, 92(3):401–419, 2002.
- [4] Ivo Babuška and Manil Suri. Locking effects in the finite element approximation of elasticity problems. *Numer. Math.*, 62(4):439–463, 1992.
- [5] Ivo Babuška and Manil Suri. On locking and robustness in the finite element method. *SIAM J. Numer. Anal.*, 29(5):1261–1293, 1992.
- [6] Susanne C. Brenner. A nonconforming mixed multigrid method for the pure traction problem in planar linear elasticity. *Math. Comp.*, 63(208):435–460, S1–S5, 1994.
- [7] Susanne C. Brenner and L. Ridgway Scott. *The mathematical theory of finite element methods*, volume 15 of *Texts in Applied Mathematics*. Springer-Verlag, New York, 1994.
- [8] Susanne C. Brenner and Li-Yeng Sung. Linear finite element methods for planar linear elasticity. *Math. Comp.*, 59(200):321–338, 1992.
- [9] Z. Cai, C.-O. Lee, T. A. Manteuffel, and S. F. McCormick. First-order system least squares for linear elasticity: numerical results. *SIAM J. Sci. Comput.*, 21(5):1706–1727 (electronic), 2000. Iterative methods for solving systems of algebraic equations (Copper Mountain, CO, 1998).
- [10] Zhiqiang Cai, Thomas A. Manteuffel, Stephen F. McCormick, and Seymour V. Parter. First-order system least squares (FOSLS) for planar linear elasticity: pure traction problem. *SIAM J. Numer. Anal.*, 35(1):320–335 (electronic), 1998.
- [11] D. Cho and L. Zikatanov. A multilevel preconditioning for generalized finite element method problems on unstructured simplicial meshes. *J. Numer. Math.*, 15(3):163–180, 2007.
- [12] Richard S. Falk. Nonconforming finite element methods for the equations of linear elasticity. *Math. Comp.*, 57(196):529–550, 1991.
- [13] Leopoldo P. Franca and Rolf Stenberg. Error analysis of Galerkin least squares methods for the elasticity equations. *SIAM J. Numer. Anal.*, 28(6):1680–1697, 1991.
- [14] Chang-Ock Lee. A conforming mixed finite element method for the pure traction problem of linear elasticity. *Appl. Math. Comput.*, 93(1):11–29, 1998.
- [15] S.V. Nepomnyaschikh. Decomposition and fictitious domains methods for elliptic boundary value problems. *SIAM, Philadelphia*, pages 581–613, 1992. Fifth International Symposium on Domain Decomposition Methods for Partial Differential Equations.
- [16] Yousef Saad. *Iterative methods for sparse linear systems*. Society for Industrial and Applied Mathematics, Philadelphia, PA, second edition, 2003.
- [17] Joachim Schöberl. Multigrid methods for a parameter dependent problem in primal variables. *Numer. Math.*, 84(1):97–119, 1999.
- [18] J. Xu. The auxiliary space method and optimal multigrid preconditioning techniques for unstructured grids. *Computing*, 56(3):215–235, 1996. International GAMM-Workshop on Multi-level Methods (Meisendorf, 1994).
- [19] Jinchao Xu. Iterative methods by space decomposition and subspace correction. *SIAM Rev.*, 34(4):581–613, 1992.

DEPARTMENT OF MATHEMATICS, THE PENNSYLVANIA STATE UNIVERSITY, UNIVERSITY PARK, PA 16802, USA.

E-mail address: brannick@math.psu.edu

DEPARTMENT OF MATHEMATICS, THE PENNSYLVANIA STATE UNIVERSITY, UNIVERSITY PARK, PA 16802, USA.

E-mail address: cho@math.psu.edu